# Fasting before your surgery or procedure – Adults & young people 16+ years



Patient Information

Anaesthesia & Pain Management

#### **Eating & Drinking**

It is important for your safety to follow these eating & drinking instructions carefully - otherwise your surgery or procedure could be delayed or cancelled.

It is important to stop eating and drinking at the correct times. *Early enough to allow your stomach to be empty but late enough to keep you hydrated.* 

### If you are due for your surgery or procedure on a <u>morning</u> or <u>all day</u> list

- Stop all food (including milk) from 2am (the night before)
- After 2am, continue with just water until
  6am. If still awake it is safe to drink one glass or cup of water per hour. An easy idea is to have a glass or cup of water after you wake up, and have it finished by 6am
- Then nothing at all by mouth after 6am no liquids, chewing gum, sweets or lollies.

## If you are due for your surgery or procedure on an afternoon list

- You can have a light breakfast (such as toast, cereal, fruit, milk, tea, coffee) the morning of your surgery or procedure, but it must be finished by 6.30am. Not fatty or fried food.
- After 6.30am, continue with just water until 10.30am. It is safe to drink one glass or cup of water per hour. Have your last glass of water finished by 10.30am.
- Then nothing at all by mouth after 10.30am
  no liquids, chewing gum, sweets or lollies.

In some specific cases for your benefit – medical or nursing staff may decide to change the times for, amounts of or types of drinks that you can have.

#### **Medicines**

It is important for your safety to follow these instructions carefully - otherwise your surgery or procedure could be delayed or cancelled.

Take all your routine medicines by mouth during the time you are allowed to drink water (unless told otherwise – see below).

If you find you have to take a medicine (or are asked by medical or nursing staff to take a medicine) **less than 2 hours** before your surgery or procedure, it is safe to take sips of water to help you swallow tablets.

In many cases, medical or nursing staff will give

nstructions	<u> </u>			
you hav	e any qu	estions <sub> </sub>	olease co	ntac