

Laparoscopy

Patient Information

Women's Health Service

Surname: NHI:

First Names:

Date of Birth: / / Sex:

PLACE PATIENT ID HERE



Going Home

You will usually be able to go home within 24 hours of your operation. You will need to arrange for someone to drive you home. You should have someone to stay with you for the first 24 hours.

You may feel drowsy because a small amount of the anaesthetic may still remain in your body. This can slow your reaction time, coordination and affect your judgement for the next 24 hours. You should not drink alcohol, make any important decisions, sign any documents (including legal), drive a car, cycle, ride a motorbike or use heavy machinery for at least 24 hours after your operation.

Some side effects following anaesthesia may include; sore/dry throat, or a slight cough. This usually resolves within a few days. Nausea, discomfort and tiredness are common side effects during the first few days following surgery.

Pain relief

Take pain relief as instructed.

You may feel some chest, shoulder or abdominal discomfort for a few days as a result of the gas introduced into your abdomen during surgery. Your body will absorb this gas within 24-48 hours which will relieve these symptoms. It can be helped by moving/walking around, taking pain relief such as Paracetamol (Panadol) and/or antiinflammatories such as ibuprofen. A heat-pack may also be helpful.

If pain becomes worse, please contact Women's Health Acute Assessment Unit on (04) 806 0740.

If you have difficulty breathing, please ring 111 or emergency assistance.

Wound Care

Each incision (cut) usually has 1 or 2 stitches. The stitches usually dissolve (disappear) within 5-7 days. If they have not dissolved after 7 days you should contact your GP and make an appointment for the practice nurse to remove them.

If you have any dressings or steri-strips on the incisions they may be removed after 48-72 hours.

Sometimes the cuts may be held together with surgical glue. If your wounds have been glued you will not need to worry about stitches and the glue will slowly peel off as the wounds heal.

You can shower as normal, pat the incisions dry with a clean towel – it is important to keep these as clean and dry as possible. Do not put creams, powders or perfume near your wound while it is still healing as these can irritate the skin.

Your wound may feel itchy, tingly and slightly lumpy. Please do not scratch or pull off the scab as these help your wound to heal.

Activity and Exercise

You will need to take it easy for a few days. You will usually be able to return to work within 1-2 weeks, but this depends on the exact treatment you have. A full recovery can take 1-2 weeks. It is important for your recovery that you have some gentle exercise each day. More strenuous exercise should be restricted for 2 weeks.

You may start driving when you are able to move from side to side, can do an emergency stop without any pain and you are no longer taking any strong pain killers.

You can start sexual activity again after 2-4 weeks and when vaginal spotting has stopped.

[Continued]

Bowel Care

Constipation (not being able to poo) can be a problem for some people after an anaesthetic. Include plenty of fresh fruit, vegetables and at least 8 glasses of water a day in your diet to keep your bowel habits regular.

Alpine tea and Kiwicrush may be helpful.

If you are have problems with constipation, please check that your discharge medication includes laxatives.

Vaginal Bleeding

You may have some light vaginal bleeding, spotting or brown discharge up to 4-6 weeks after surgery, this is normal as long as it is not unpleasant smelling.

If it is unpleasant smelling, this could be because of an infection and you may need to be treated with antibiotics.

If you have heavy, bright red bleeding and you are filling one sanitary pad every hour, please contact the Women's Health Acute Assessment Unit, or your local Emergency Department.

It is important to use pads instead of tampons and AVOID swimming or hot tub soaks until you have stopped bleeding to avoid infection.

Complications to look out for

We particularly want you to tell us if:

- You have increasing pain not controlled by pain relief.
- You have a fever (temperature).
- Your wound is bleeding and will not stop.
- Your wound is red or warm to the touch.
- You have thick and/or smelly ooze from your wound.
- You have heavy vaginal bleeding or offensive smelling discharge.

- You are constipated or unable to pass urine.
- You have on-going nausea (feeling sick) or vomiting.
- You notice one or both of your calf/s has become sore to the touch, is red or there is an increase in size of your calf/s.

Discharge and Follow-up

Following your operation the doctor will tell you of their findings. Your discharge summary is provided to you and your GP. Please check your discharge summary for details of your follow-up appointment.

At this time your recovery will be assessed and any results will be discussed.

Most women do not have any problems following Laparoscopic surgery.

If you have any concerns regarding your post operative recovery, please contact the **Gynaecology Ward/Womens Health Assessment Unit** on **04 806 0740 (24 hour coverage)**.

Contact number if you have not heard about lab results and if you need a follow up appointment:

**Women's Clinics Reception
Level 3
Wellington Regional Hospital
Phone: 04 806 0760**

Reception will either take a message or forward to the Clinic Nurse for the Specialist whose care you are assigned.